

NUHEP-TH-92-23

UCD-92-25

February 1993

(revised version)

Gluon Fragmentation into Heavy Quarkonium

Eric Braaten

Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208

Tzu Chiang Yuan

Davis Institute for High Energy Physics

Department of Physics, University of California, Davis, CA 95616

Abstract

The dominant production mechanism for heavy quark-antiquark bound states in very high energy processes is fragmentation, the splitting of a high energy parton into a quarkonium state and other partons. We show that the fragmentation functions $D(z, \mu)$ describing these processes can be calculated using perturbative QCD. We calculate the fragmentation functions for a gluon to split into S-wave quarkonium states to leading order in the QCD coupling constant. The leading logarithms of μ/m_Q , where μ is the factorization scale and m_Q is the heavy quark mass, are summed up using Altarelli-Parisi evolution equations.

Quantitative evidence for quantum chromodynamics (QCD) as the fundamental field theory describing the strong interactions has come primarily from high energy processes involving leptons and the electroweak gauge bosons. Such processes are simpler than most purely hadronic processes, because leptons and electroweak gauge bosons do not have strong interactions. The next simplest particles as far as the strong interactions are concerned are heavy quarkonia, the bound states of a heavy quark and antiquark. While not pointlike, the lowest states in the charmonium and bottomonium systems have typical radii that are significantly smaller than those of hadrons containing light quarks. They have simple internal structure, consisting primarily of a nonrelativistic quark and antiquark only. The charmonium and bottomonium systems exhibit a rich spectrum of orbital and angular excitations. Thus in addition to being simple enough to be used as probes of the strong interactions, heavy quarkonia are also a potentially much richer source of information than leptons and electroweak gauge bosons.

In most previous studies of the production of heavy quarkonia in high energy processes, it was implicitly assumed that they are produced by *short distance* mechanisms, in which the heavy quark and antiquark are created with transverse separations of order $1/E$, where E is the characteristic energy scale of the process. In this paper, we point out that the dominant mechanism at very high energies is *fragmentation*, the production of a high energy parton followed by its splitting into the quarkonium state and other partons. The $Q\bar{Q}$ pair are created with a separation of order $1/m_Q$, where m_Q is the mass of the heavy quark Q . The fragmentation mechanism is often of higher order in the QCD coupling constant α_s than the short distance mechanism, but it is enhanced by a factor of $(E/m_Q)^2$ and thus dominates at high energies $E \gg m_Q$. The fragmentation of a parton into a quarkonium state is described by a fragmentation function $D(z, \mu)$, where z is the longitudinal momentum fraction of the quarkonium state and μ is a factorization scale. We calculate to leading order in α_s the fragmentation functions $D(z, m_Q)$ for gluons to split into S-wave quarkonium states at energy scales μ of order m_Q . The fragmentation functions at larger scales μ are then determined by Altarelli-Parisi evolution equations which sum up the leading logarithms of μ/m_Q .

One of the quarkonium processes that is important in hadron collider physics is the

production of charmonium at large transverse momentum p_T . A charmonium state can either be produced directly at large p_T or it can be produced indirectly by the decay of a large p_T B -meson or a higher charmonium state with large p_T . In previous calculations of the rate for direct production of charmonium at large p_T [1], the dominant mechanisms were assumed to be short distance processes, in which a collinear $c\bar{c}$ pair in a color-singlet S-wave state is created with transverse separation on the order of $1/p_T$. A typical Feynman diagram which contributes to the production of the 1S_0 charmonium state η_c at order α_s^3 is the diagram for $gg \rightarrow c\bar{c}g$ shown in Figure 1. The order- α_s^4 radiative corrections to this process include the Feynman diagram for $gg \rightarrow c\bar{c}gg$ shown in Figure 2. In most regions of phase space, the virtual gluons in Figure 2 are off their mass shells by amounts of order p_T , and the contribution from this diagram is suppressed relative to the diagram in Figure 1 by a power of the running coupling constant $\alpha_s(p_T)$. But there is a part of the phase space in which the virtual gluon attached to the $c\bar{c}$ pair in Figure 2 is off-shell by an amount of order m_c . The propagator of this virtual gluon enhances the cross section by a factor of p_T^2/m_c^2 . At large enough p_T , this easily overwhelms the extra power of the coupling constant α_s . The enhancement is due to the fact that the $c\bar{c}$ pair can be produced with transverse separation of order $1/m_c$ instead of $1/p_T$.

A more thorough analysis of the amplitude for $gg \rightarrow \eta_c gg$ reveals that the term that is enhanced by p_T^2/m_c^2 can be written in a factored form. The first factor is the amplitude for the production of a virtual gluon g^* with high- p_T but low invariant mass q^2 via the process $gg \rightarrow gg^*$. In the limit $q^2 \ll p_T^2$, it reduces to the on-shell scattering amplitude for $gg \rightarrow gg$. The second factor is the propagator $1/q^2$ for the virtual gluon. The third and final factor is the amplitude for the process $g^* \rightarrow \eta_c g$, in which an off-shell gluon fragments into an η_c and a gluon. The factoring of the amplitude allows the fragmentation contribution to the differential cross section $d\sigma_{\eta_c}(E)$ for producing an η_c with energy $E \gg m_c$ to be written in a factorized form:

$$d\sigma_{\eta_c}(E) \approx \int_0^1 dz \, d\hat{\sigma}_g(E/z) \, D_{g \rightarrow \eta_c}(z, m_c) , \quad (1)$$

where $d\hat{\sigma}_g(E)$ is the differential cross section for producing a real gluon of energy E . All of the dependence on the energy E appears in the subprocess cross section $d\hat{\sigma}_g$, while all

the dependence on the quark mass m_c is in the fragmentation function $D(z, m_c)$. The variable z is the longitudinal momentum fraction of the η_c relative to the gluon. The physical interpretation of (1) is that an η_c of energy E can be produced by first producing a gluon of larger energy E/z which subsequently splits into an η_c carrying a fraction z of the gluon energy.

The generalization of the leading order formula (1) to all orders in α_s is straightforward. At higher orders in α_s , the gluon that splits into a quarkonium state $\mathcal{O}$ can itself arise from the splitting of a higher energy parton into a collinear gluon. This splitting process gives rise to logarithms of E/m_Q . In order to maintain the factorization of the dependences on E and m_Q , it is necessary to introduce a factorization scale μ : $\log(E/m_Q) = \log(E/\mu) + \log(\mu/m_Q)$. To all orders in α_s , the fragmentation contribution to the differential cross section for producing a quarkonium state $\mathcal{O}$ with energy E can be written in the factorized form

$$d\sigma_{\mathcal{O}}(E) = \sum_i \int_0^1 dz d\hat{\sigma}_i(E/z, \mu) D_{i \rightarrow \mathcal{O}}(z, \mu), \quad (2)$$

where the sum is over all parton types i . The scale μ is arbitrary, but large logarithms of E/μ in the parton cross section $d\hat{\sigma}_i$ can be avoided by choosing μ on the order of E . Large logarithms of μ/m_Q then necessarily appear in the fragmentation functions $D_{i \rightarrow \mathcal{O}}(z, \mu)$, but they can be summed up by solving the evolution equations [2]

$$\mu \frac{\partial}{\partial \mu} D_{i \rightarrow \mathcal{O}}(z, \mu) = \sum_j \int_z^1 \frac{dy}{y} P_{i \rightarrow j}(z/y, \mu) D_{j \rightarrow \mathcal{O}}(y, \mu), \quad (3)$$

where $P_{i \rightarrow j}(x, \mu)$ is the Altarelli-Parisi function for the splitting of the parton of type i into a parton of type j with longitudinal momentum fraction x . For many applications, calculations to leading order in α_s require only the $g \rightarrow g$ splitting function, which is

$$P_{g \rightarrow g}(x, \mu) = \frac{3\alpha_s(\mu)}{\pi} \left(\frac{1-x}{x} + \frac{x}{(1-x)_+} + x(1-x) + \frac{33-2n_f}{36} \delta(1-x) \right), \quad (4)$$

where n_f is the number of light quark flavors. The boundary condition on this evolution equation is the fragmentation function $D_{i \rightarrow \mathcal{O}}(z, m_Q)$ at the scale m_Q . It can be calculated perturbatively as a series in $\alpha_s(m_Q)$.

We proceed to calculate the fragmentation function $D_{g \rightarrow \eta_c}(z, m_c)$ for a gluon to split into the 1S_0 charmonium state η_c to leading order in $\alpha_s(m_c)$. A process (such as $gg \rightarrow gg$) that produces a real gluon of 4-momentum q has a matrix element of the form $\mathcal{M}_\alpha \epsilon^\alpha(q)$, where $\epsilon^\alpha(q)$ is the polarization 4-vector of the on-shell ($q^2 = 0$) gluon. In the corresponding fragmentation process (such as Figure 2), a virtual gluon is produced with large energy $q_0 \gg m_c$ but small invariant mass $s = q^2$ of order m_c^2 , and it subsequently fragments into an η_c and a real gluon. The fragmentation probability $\int_0^1 dz D(z, m_c)$ is the ratio of the rates for these two processes. In Feynman gauge, the fragmentation term in the matrix element for the η_c production has the form $\mathcal{M}_\alpha (-ig^{\alpha\beta}/q^2) \mathcal{A}_\beta$, where $\mathcal{M}_\alpha$ is the matrix element for the production of the virtual gluon and $\mathcal{A}_\beta$ is the amplitude for $g^* \rightarrow \eta_c g$. The fragmentation term is distinguished from the short distance terms in the matrix element by the presence of the small denominator q^2 of order m_c^2 . The amplitude $\mathcal{A}_\beta$ can be written down using standard Feynman rules for quarkonium processes [3]. Multiplying $\mathcal{A}_\alpha$ by its complex conjugate and summing over final colors and spins, we get

$$\begin{aligned} \sum \mathcal{A}_\alpha \mathcal{A}_\beta^* = & \frac{16\pi}{3} \alpha_s^2 \frac{|R(0)|^2}{2m_c} \frac{1}{(s - 4m_c^2)^2} \left(- (s - 4m_c^2)^2 g_{\alpha\beta} \right. \\ & \left. + 2(s + 4m_c^2)(p_\alpha q_\beta + q_\alpha p_\beta) - 4s p_\alpha p_\beta - 16m_c^2 q_\alpha q_\beta \right), \end{aligned} \quad (5)$$

where p is the 4-momentum of the η_c . Terms proportional to q_α or q_β are gauge artifacts and can be dropped. In an appropriate axial gauge, q_α and q_β are of order m_c^2/q_0 when contracted with the numerator of the propagator of the virtual gluon. In covariant gauges, the q_α and q_β terms are not suppressed but are cancelled by other diagrams. In the $p_\alpha p_\beta$ term, we can set $p = zq + p_\perp$ up to corrections of order m_c^2/q_0 , where z is the longitudinal momentum fraction and $p_\perp$ is the transverse part of the 4-vector p . In a frame where $q = (q_0, 0, 0, q_3)$, $z = (p_0 + p_3)/(q_0 + q_3)$ and $p_\perp = (0, p_1, p_2, 0)$. After averaging over the directions of the transverse momentum, $p_\perp^\alpha p_\perp^\beta$ can be replaced by $-g^{\alpha\beta} \vec{p}_\perp^2/2$, up to terms that are suppressed in axial gauge. The terms in (5) that contribute to fragmentation then reduce to

$$\sum \mathcal{A}_\alpha \mathcal{A}_\beta^* = \frac{16\pi}{3} \alpha_s^2 \frac{|R(0)|^2}{2m_c} \frac{1}{(s - 4m_c^2)^2} \left((s - 4m_c^2)^2 - 2(1 - z)(zs - 4m_c^2)s \right) (-g_{\alpha\beta}). \quad (6)$$

We have used the conservation of the $q_0 - q_3$ component of the 4-momentum in the form $s = (\vec{p}_\perp^2 + 4m_c^2)/z + \vec{p}_\perp^2/(1-z)$. At this point, it is easy to calculate the rate for production of $\eta_c g$ in the limit $q_0^2 \gg q^2 \sim m_c^2$ and divide it by the rate for production of an on-shell gluon. The resulting fragmentation probability is

$$\int_0^1 dz D_{g \rightarrow \eta_c}(z) = \frac{\alpha_s^2}{3\pi} \frac{|R(0)|^2}{2m_c} \int_{4m_c^2}^\infty ds \int_{4m_c^2/s}^1 dz \frac{s^2 + 16m_c^4 - 2z(s + 4m_c^2)s + 2z^2 s^2}{s^2(s - 4m_c^2)^2}, \quad (7)$$

where $R(0)$ is the nonrelativistic radial wavefunction at the origin for the S-wave bound state. We have increased the upper endpoint of the integration over s to infinity, because the resulting error is of order m_c^2/q_0^2 , which we have been consistently neglecting. Interchanging orders of integration, we can read off the fragmentation function:

$$D_{g \rightarrow \eta_c}(z, 2m_c) = \frac{1}{3\pi} \alpha_s(2m_c)^2 \frac{|R(0)|^2}{M_{\eta_c}^3} \left(3z - 2z^2 + 2(1-z) \log(1-z) \right). \quad (8)$$

We have set the scale in the fragmentation function and in the running coupling constant to $\mu = 2m_c$, which is the minimum value of the invariant mass $\sqrt{s}$ of the fragmenting gluon. In the denominator, we have set $2m_c = M_{\eta_c}$, which takes into account the correct phase space limitations and is accurate up to relativistic corrections. The value of the S-state wavefunction at the origin $R(0)$ is determined from the ψ electronic width to be $|R(0)|^2 = (0.8 \text{ GeV})^3$. We use the value $\alpha_s(2m_c) = 0.26$ for the strong coupling constant.

Given the initial fragmentation function (8), the fragmentation function is determined at larger values of μ by solving the evolution equation (3) with (8) as a boundary condition. The z -dependence of $D_{g \rightarrow \eta_c}(z, \mu)$ at the energy scales $\mu = 2m_c$ and $\mu = 20m_c$ is illustrated in Figure 3. The evolution causes the fragmentation function to decrease at large z and to diverge at $z = 0$. A physical cross section like (2) will still be well-behaved, because phase space limitations will place an upper bound on the parton energy E/z which translates into a lower bound on z . It is evident from Figure 3 that taking into account the evolution of the fragmentation function can significantly increase the rate for the production process, particularly at small values of z .

The fragmentation function for a gluon into J/ψ can be calculated to leading order in α_s from the Feynman diagrams for $g^* \rightarrow \psi g g$. The square of the amplitude $\sum \mathcal{A}_\alpha \mathcal{A}_\beta^*$ for

this process can be extracted from a calculation of the matrix element for $e^+e^- \rightarrow \psi gg$ [4]. The calculation of the fragmentation function is rather involved and we present only the final result:

$$D_{g \rightarrow \psi}(z, 2m_c) = \frac{5}{144\pi^2} \alpha_s(2m_c)^3 \frac{|R(0)|^2}{M_\psi^3} \int_0^z dr \int_{(r+z^2)/2z}^{(1+r)/2} dy \frac{1}{(1-y)^2(y-r)^2(y^2-r)^2} \sum_{i=0}^2 z^i \left(f_i(r, y) + g_i(r, y) \frac{1+r-2y}{2(y-r)\sqrt{y^2-r}} \log \frac{y-r+\sqrt{y^2-r}}{y-r-\sqrt{y^2-r}} \right). \quad (9)$$

where the integration variables are $r = 4m_c^2/s$ and $y = p \cdot q/s$. The functions f_i and g_i are

$$\begin{aligned} f_0(r, y) &= r^2(1+r)(3+12r+13r^2) - 16r^2(1+r)(1+3r)y \\ &- 2r(3-9r-21r^2+7r^3)y^2 + 8r(4+3r+3r^2)y^3 - 4r(9-3r-4r^2)y^4 \\ &- 16(1+3r+3r^2)y^5 + 8(6+7r)y^6 - 32y^7, \end{aligned} \quad (10)$$

$$\begin{aligned} f_1(r, y) &= -2r(1+5r+19r^2+7r^3)y + 96r^2(1+r)y^2 + 8(1-5r-22r^2-2r^3)y^3 \\ &+ 16r(7+3r)y^4 - 8(5+7r)y^5 + 32y^6, \end{aligned} \quad (11)$$

$$\begin{aligned} f_2(r, y) &= r(1+5r+19r^2+7r^3) - 48r^2(1+r)y - 4(1-5r-22r^2-2r^3)y^2 \\ &- 8r(7+3r)y^3 + 4(5+7r)y^4 - 16y^5, \end{aligned} \quad (12)$$

$$\begin{aligned} g_0(r, y) &= r^3(1-r)(3+24r+13r^2) - 4r^3(7-3r-12r^2)y - 2r^3(17+22r-7r^2)y^2 \\ &+ 4r^2(13+5r-6r^2)y^3 - 8r(1+2r+5r^2+2r^3)y^4 - 8r(3-11r-6r^2)y^5 \\ &+ 8(1-2r-5r^2)y^6, \end{aligned} \quad (13)$$

$$\begin{aligned} g_1(r, y) &= -2r^2(1+r)(1-r)(1+7r)y + 8r^2(1+3r)(1-4r)y^2 \\ &+ 4r(1+10r+57r^2+4r^3)y^3 - 8r(1+29r+6r^2)y^4 - 8(1-8r-5r^2)y^5, \end{aligned} \quad (14)$$

$$\begin{aligned} g_2(r, y) &= r^2(1+r)(1-r)(1+7r) - 4r^2(1+3r)(1-4r)y \\ &- 2r(1+10r+57r^2+4r^3)y^2 + 4r(1+29r+6r^2)y^3 + 4(1-8r-5r^2)y^4. \end{aligned} \quad (15)$$

The integrals over r and y in (9) must be evaluated numerically to obtain the fragmentation function at the energy scale $\mu = 2m_c$. At larger values of μ , it is found by solving the evolution equation (3) with (9) as a boundary condition. The z -dependence of $D_{g \rightarrow \psi}(z, \mu)$ at the energy scales $\mu = 2m_c$ and $\mu = 20m_c$ is illustrated in Figure 4. The evolution causes the fragmentation function to decrease at large z and to diverge at $z = 0$.

An order of magnitude estimate of the gluon fragmentation contribution to quarkonium production in any high energy process can be obtained by multiplying the cross section for producing gluons of energy $E > 2m_Q$ by the initial fragmentation probability $\int_0^1 dz D(z, 2m_Q)$. For the η_c and ψ , these probabilities are $4.6 \cdot 10^{-5}$ and $2.8 \cdot 10^{-6}$. Thus, we can expect the asymptotic production rate of ψ to be more than an order of magnitude smaller than η_c . The initial fragmentation function for the splitting of a gluon into the 3S_1 bottomonium state Υ is also given by (9), except that the mass M_ψ is replaced by M_Υ , the scale $2m_c$ is replaced by $2m_b$, and $R(0)$ is the appropriate wavefunction at the origin. The initial fragmentation probability is about $5.3 \cdot 10^{-7}$, which is smaller than for ψ by about a factor of 5.

In hadron colliders, short distance processes dominate the direct production of charmonium at small p_T , because fragmentation processes are suppressed by powers of $\alpha_s(m_c)$. At sufficiently large p_T , fragmentation processes must dominate, because the short distance processes are suppressed by powers of m_c^2/p_T^2 . We can make a quantitative estimate of the p_T at which the crossover occurs by comparing the differential cross sections for short distance process with the differential cross section for the gluon scattering process $gg \rightarrow gg$ multiplied by the appropriate fragmentation probability. For simplicity, we consider 90° scattering in the gg center of mass frame. In terms of p_T , we have $s = 4p_T^2$. In the limit $p_T \gg m_c$, the differential cross section for $gg \rightarrow g\eta_c$ is [1] $d\sigma/dt = 81\pi\alpha_s^3|R(0)|^2/(256M_{\eta_c}p_T^6)$. The differential cross section for $gg \rightarrow gg$ is $d\sigma/dt = 243\pi\alpha_s^2/(128p_T^4)$. To allow for fragmentation of either of the two outgoing gluons, we multiply by twice the initial fragmentation probability $\alpha_s^2|R(0)|^2/(9\pi M_{\eta_c}^3)$. The resulting differential cross section exceeds that for the short distance process at $p_T \approx \sqrt{3\pi/4\alpha_s}M_{\eta_c} \approx 3M_{\eta_c}$. The differential cross section for $gg \rightarrow g\psi$ is [1] $d\sigma/dt = 5\pi\alpha_s^3|R(0)|^2M_\psi/(128p_T^8)$ and for this case we estimate the crossover point to be $p_T \approx \sqrt{1.05/\alpha_s}M_\psi \approx 2M_\psi$. Thus fragmentation should dominate over short distance production at values of p_T that are being measured in present collider experiments.

The importance of fragmentation for charmonium production in high energy processes can also be seen from the surprisingly large rate [5] for $Z^0 \rightarrow \psi c\bar{c}$, which is two orders of magnitude larger than that of $Z^0 \rightarrow \psi gg$. The explanation for this is that $Z^0 \rightarrow \psi gg$ is

a short distance process, while $Z^0 \rightarrow \psi c\bar{c}$ includes a fragmentation contribution enhanced by $(M_Z/M_\psi)^2$. The enhanced contribution arises from the decay $Z^0 \rightarrow c\bar{c}$, followed by the splitting $c \rightarrow \psi c$ or $\bar{c} \rightarrow \psi\bar{c}$. With this insight, the lengthy calculation presented in Ref. [5] can be reduced to a simple calculation of the fragmentation function $D_{c \rightarrow \psi}(z, M_Z)$. This calculation will be presented elsewhere [6].

The probability for a virtual gluon to decay into a ψ was calculated by Hagiwara, Martin and Stirling [7] and used to study J/ψ production from gluon jets at the LEP collider. They did not calculate the fragmentation function $D_{g \rightarrow \psi}(z, \mu)$ and thus were unable to sum up large logarithms of M_Z/m_c . Furthermore, their expression for the fragmentation probability is missing a factor of $1/(16\pi^2)$, which explains the surprisingly large rate that they found for this ψ -production mechanism.

A complete calculation of the fragmentation contribution to ψ production in high energy processes must include the production of the P-wave charmonium states χ_{cJ} , followed by their radiative decays into ψ . In calculating the fragmentation functions for gluons to split into P-wave charmonium states, there are two distinct contributions that must be included at leading order in α_s . The P-wave state can arise either from the production of a collinear $c\bar{c}$ pair in a color-singlet P-wave state, or from the production of a collinear $c\bar{c}$ pair in a color-octet S-wave state [8]. Calculations of the P-wave fragmentation functions will be presented elsewhere [9].

We have shown in this letter that the dominant production mechanism for quarkonium in high energy processes is fragmentation, the production of a high energy parton followed by its splitting into the quarkonium state. We calculated the fragmentation functions $D(z, \mu)$ for gluons to split into S-wave quarkonium states to leading order in α_s . The fragmentation functions satisfy Altarelli-Parisi evolution equations which can be used to sum up large logarithms of μ/m_Q . Most previous calculations of quarkonium production have considered only short-distance production mechanisms, because the fragmentation mechanism is often of higher order in α_s . At high energies E , the fragmentation mechanism dominates because it is enhanced by a factor of $(E/m_Q)^2$. In the case of charmonium production at large p_T in hadron colliders, we estimated the transverse momentum at which fragmentation begins to

dominate to be less than 10 GeV. All previous calculations of quarkonium production from this and other high energy processes must therefore be reexamined, taking into account the possibility of production by fragmentation.

This work was supported in part by the U.S. Department of Energy, Division of High Energy Physics, under Grant DE-FG02-91-ER40684. We would like to acknowledge G.P. Lepage for suggesting the importance of fragmentation in the production of quarkonium in high energy processes. We thank M.L. Mangano for pointing out an error in an earlier version of this paper. We would also like to acknowledge useful conversations with G.T. Bodwin, K. Cheung, J. Gunion, and W.J. Stirling.

References

- [1] See, for example, R. Baier and R. Rückl, *Z. Phys.* **C19**, 251 (1983).
- [2] R.D. Field, *Applications of Perturbative QCD*, (Addison Wesley, 1989).
- [3] J.H. Kühn, J. Kaplan, and E.G.O. Safiani, *Nucl. Phys.* **B157**, 125 (1979); B. Guberina, J.H. Kühn, R.D. Peccei, and R. Rückl, *Nucl. Phys.* **B174**, 317 (1980).
- [4] J.H. Kühn and H. Schneider, *Phys. Rev.* **D24**, 2996 (1981); *Z. Phys.* **C11**, 263 (1981).
- [5] V. Barger, K. Cheung, and W.-Y. Keung, *Phys. Rev.* **D41**, 1541 (1990).
- [6] E. Braaten, K. Cheung, and T.C. Yuan, Northwestern preprint NUHEP-TH-93-2 (February, 1993).
- [7] K. Hagiwara, A.D. Martin, and W.J. Stirling, *Phys. Lett.* **B267**, 527 (1991) and erratum (to be published).
- [8] G.T. Bodwin, E. Braaten, T.C. Yuan, and G.P. Lepage, *Phys. Rev.* **D46**, R3703 (1992).
- [9] E. Braaten and T.C. Yuan (in preparation).

Figure Captions

1. A Feynman diagram for $gg \rightarrow c\bar{c}g$ that contributes to η_c production at order α_s^3 .
2. A Feynman diagram for $gg \rightarrow c\bar{c}gg$ that contributes to η_c production at order α_s^4 .
3. The fragmentation function $D_{g \rightarrow \eta_c}(z, \mu)$ as a function of z for $\mu = 2m_c$ (solid line) and $\mu = 20m_c$ (dotted line).
4. The fragmentation function $D_{g \rightarrow \psi}(z, \mu)$ as a function of z for $\mu = 2m_c$ (solid line) and $\mu = 20m_c$ (dotted line).